

# Team Round

Lexington High School

April 9, 2016

## [70] Potpourri

1. Suppose that 20% of a number is 17. Find 20% of 17% of the number.
2. Let  $A, B, C, D$  represent the numbers 1 through 4 in some order, with  $A \neq 1$ . Find the maximum possible value of

$$\frac{\log_A B}{C + D}.$$

Here,  $\log_A B$  is the unique real number  $X$  such that  $A^X = B$ .

3. There are six points in a plane, no four of which are collinear. A line is formed connecting every pair of points. Find the smallest possible number of distinct lines formed.
4. Let  $a, b, c$  be real numbers which satisfy

$$\begin{aligned}\frac{2017}{a} &= a(b + c) \\ \frac{2017}{b} &= b(a + c) \\ \frac{2017}{c} &= c(a + b).\end{aligned}$$

Find the sum of all possible values of  $abc$ .

5. Let  $a$  and  $b$  be complex numbers such that  $ab + a + b = (a + b + 1)(a + b + 3)$ . Find all possible values of  $\frac{a+1}{b+1}$ .
6. Let  $\triangle ABC$  be a triangle. Let  $X, Y, Z$  be points on lines  $BC, CA$ , and  $AB$ , respectively, such that  $X$  lies on segment  $BC$ ,  $B$  lies on segment  $AY$ , and  $C$  lies on segment  $AZ$ . Suppose that the circumcircle of  $\triangle XYZ$  is tangent to lines  $AB, BC$ , and  $CA$  with center  $I_A$ . If  $AB = 20$  and  $I_A C = AC = 17$  then compute the length of segment  $BC$ .
7. An ant makes 4034 moves on a coordinate plane, beginning at the point  $(0, 0)$  and ending at  $(2017, 2017)$ . Each move consists of moving one unit in a direction parallel to one of the axes. Suppose that the ant stays within the region  $|x - y| \leq 2$ . Let  $N$  be the number of paths the ant can take. Find the remainder when  $N$  is divided by 1000.
8. A 10 digit positive integer  $\overline{a_9 a_8 a_7 \cdots a_1 a_0}$  with  $a_9$  nonzero is called *deceptive* if there exist distinct indices  $i > j$  such that  $\overline{a_i a_j} = 37$ . Find the number of deceptive positive integers.
9. A circle passing through the points  $(2, 0)$  and  $(1, 7)$  is tangent to the  $y$ -axis at  $(0, r)$ . Find all possible values of  $r$ .
10. An ellipse with major and minor axes 20 and 17, respectively, is inscribed in a square whose diagonals coincide with the axes of the ellipse. Find the area of the square.

**[130] Long Answer**

1. [10] Find with proof the smallest positive integer  $k$  such that every  $k$ -element subset of  $\{1, 2, 3, \dots, 500\}$  contains two distinct elements  $a, b$  such that  $a + b$  is also an element of the set.
2. [15] Let  $\alpha = \frac{\sqrt{5}+1}{2}$ . Find all ordered pairs of positive integers  $(m, n)$  with  $m \neq n$  such that  $\{\alpha^m\} = \{\alpha^n\}$ . (Here  $\{x\}$  denotes the fractional part of  $x$ ).
3. [30] The goal of this problem is to show that the maximum area of a polygon with a fixed number of sides and a fixed perimeter is achieved by a regular polygon.
  - (a) [4] Prove that the polygon with maximum area must be convex. (Hint: If any angle is concave, show that the polygon's area can be increased.)
  - (b) [8] Prove that if two adjacent sides have different lengths, the area of the polygon can be increased without changing the perimeter.
  - (c) [4] Prove that the polygon with maximum area is equilateral, that is, has all the same side lengths.

It is true that when given all four side lengths in order of a quadrilateral, the maximum area is achieved in the unique configuration in which the quadrilateral is cyclic, that is, it can be inscribed in a circle.

- (d) [8] Prove that in an equilateral polygon, if any two adjacent angles are different then the area of the polygon can be increased without changing the perimeter.
  - (e) [4] Prove that the polygon of maximum area must be equiangular, or have all angles equal.
  - (f) [2] Prove that the polygon of maximum area is a regular polygon.
4. [35] Let  $A$  be a list of  $N$  positive integers sorted least to greatest. Say we are searching the set for an element  $E$ . Define *trivial search* as simply searching for the element from the start of  $A$  to the end of  $A$ . This can be very inefficient for large lists.

Define *binary search* as a recursive process for sorted lists as such. Compare  $E$  to the middle element of  $A$ . If  $E$  is greater than the middle element, perform this same process on the second half of the sequence. If  $E$  is less than the middle element, perform this same process on the first half of the sequence. This continues until the middle element equals  $E$ , or the other half is empty.

For example, let  $A = \{1, 2, 3, 4, 5, 6\}$  and  $E = 4$ . We first check the range 1 to 6, where the middle element is 3. As  $4 > 3$ , we only look at the range above this middle element. This range is from 4 to 6, where the middle element is 5. As  $4 < 5$ , we only look at the range lower than this element. This range is from 4 to 4, where the middle element is 4. As this equals  $E$ , we end our search after only three comparisons.

- (a) [1] How many comparisons, at worst-case, will be needed with *binary search* for an element  $E$  on a sorted list of length 8?
- (b) [1] How many comparisons, at worst-case, will be needed with *binary search* for an element  $E$  on a sorted list of length 16?
- (c) [3] How many comparisons, at worst-case, will be needed with *binary search* for an element  $E$  on a sorted list of length  $N$ ?
- (d) [3] Prove that *binary search* will always determine whether or not  $E$  is in element in a sorted list  $A$ .
- (e) [5] Describe a method to determine the number of elements in a sorted list  $A$  that are equal to element  $E$ .

*Binary search* is a powerful tool, and can be used for a number of different problems involving searching for some quantity in an efficient manner. Binary search can even be used on the real numbers to approximate certain values.

- (a) [7] Describe a method to approximate  $\sqrt{5}$ , using binary search on the range  $[1, 5]$ .
- (b) [7] Say a sorted list  $A$  contains the first  $N$  elements of a geometric series with starting term  $a$  and ratio  $r > 1$ , but with one element removed. Describe a method to use binary search to determine the removed element if you are given  $a$  and  $r$ .
- (c) [8] Say a sorted list  $A$  has all of its elements rotated to the left by  $k$  elements (a list  $\{1, 2, 3\}$  rotated by 2 becomes  $\{2, 3, 1\}$ ). Describe a method to use binary search to determine the value of  $k$ .
5. [40] Let  $P$  be a point and  $\omega$  be a circle with center  $O$  and radius  $r$ . We define the **power** of the point  $P$  with respect to the circle  $\omega$  to be  $OP^2 - r^2$ , and we denote this by  $\text{pow}(P, \omega)$ . We define the **radical axis** of two circles  $\omega_1$  and  $\omega_2$  to be the locus of all points  $P$  such that  $\text{pow}(P, \omega_1) = \text{pow}(P, \omega_2)$ . It turns out that the pairwise radical axes of three circles are either concurrent or pairwise parallel. The concurrence point is referred to as the **radical center** of the three circles.

In  $\triangle ABC$ , let  $I$  be the incenter,  $\Gamma$  be the circumcircle, and  $O$  be the circumcenter. Let  $A_1, B_1, C_1$  be the point of tangency of the incircle of  $\triangle ABC$  with side  $BC, CA, AB$ , respectively. Let  $X_1, X_2 \in \Gamma$  such that  $X_1, B_1, C_1, X_2$  are collinear in this order. Let  $M_A$  be the midpoint of  $BC$ , and define  $\omega_A$  as the circumcircle of  $\triangle X_1 X_2 M_A$ . Define  $\omega_B, \omega_C$  analogously. The goal of this problem is to show that the radical center of  $\omega_A, \omega_B, \omega_C$  lies on line  $\overline{OI}$ .

- (a) [4] Let  $A'_1$  denote the intersection of  $B_1 C_1$  and  $BC$ . Show that  $\frac{A_1 B}{A_1 C} = \frac{A'_1 B}{A'_1 C}$ .
- (b) [8] Prove that  $A_1$  lies on  $\omega_A$ .
- (c) [6] Prove that  $A_1$  lies on the radical axis of  $\omega_B$  and  $\omega_C$ .
- (d) [13] Prove that the radical axis of  $\omega_B$  and  $\omega_C$  is perpendicular to  $B_1 C_1$ .
- (e) [2] Prove that the radical center of  $\omega_A, \omega_B, \omega_C$  is the orthocenter of  $\triangle A_1 B_1 C_1$ .
- (f) [7] Conclude that the radical center of  $\omega_A, \omega_B, \omega_C, O$ , and  $I$  are collinear.